arXiv:hep-th/9706219v2 3 Jul 1997

OU-HET 269
KEK-Prep.97-93
KEK-TH-524
hep-th/9706219

Low Energy Theorems in N=1 Supersymmetric Theory

Kiyoshi Higashijima,[1*] **Muneto Nitta,**[1,2†] **Kazutoshi Ohta**[1‡]
and
Nobuyoshi Ohta[1§]

[1] *Department of Physics, Graduate School of Science, Osaka University,*
Toyonaka, Osaka 560, Japan
[2] *Theory Division, Institute of Particle and Nuclear Studies, KEK,*
Tsukuba, Ibaraki 305, Japan

Abstract

In $N = 1$ supersymmetric theories, quasi Nambu-Goldstone (QNG) bosons appear in addition to ordinary Nambu-Goldstone (NG) bosons when the global symmetry G breaks down spontaneously. We investigate two-body scattering amplitudes of these bosons in the low-energy effective Lagrangian formalism. They are expressed by the curvature of Kähler manifold. The scattering amplitudes of QNG bosons are shown to coincide with those of NG bosons though the effective Lagrangian contains an arbitrary function, and those with odd number of QNG bosons all vanish.

---

[*] higashij@phys.wani.osaka-u.ac.jp.

[†] nittam@theory.kek.jp, nitta@phys.wani.osaka-u.ac.jp.

[‡] kohta@funpth.phys.sci.osaka-u.ac.jp. Supported in part by the JSPS Research Fellowships.

[§] ohta@phys.wani.osaka-u.ac.jp.

1 Introduction

When the global symmetry G breaks down spontaneously to its subgroup H, there are massless Nambu-Goldstone (NG) bosons, as many as the broken generators. The low energy theorems say that the low-energy scattering amplitudes of NG bosons are determined only by the symmetries G and H. If we integrate out the massive particles, we obtain the low-energy effective Lagrangian which reproduces the scattering amplitudes of NG bosons.

Callan, Coleman, Wess and Zumino (CCWZ) constructed such a low-energy effective Lagrangian as a nonlinear sigma model whose target manifold is the coset G/H parametrized by NG bosons [1]. We can calculate the low-energy scattering amplitudes of NG bosons by summing up tree graphs. The target manifold G/H is homogeneous and compact and has G-invariant metric. The broken symmetry G is realized nonlinearly while the unbroken symmetry H is represented linearly.

In theories with unbroken $N = 1$ supersymmetry, there come out some extra massless quasi Nambu-Goldstone (QNG) bosons besides the ordinary NG bosons. The target manifold of the $N = 1$ nonlinear sigma model in four dimensions must be Kähler [2] with G-isometry and is parametrized by the NG and QNG bosons. General methods to construct Kähler potential which determines the effective Lagrangian are given by Bando, Kuramoto, Maskawa and Uehara (BKMU) in their celebrated papers [3].

In supersymmetric theories, the global symmetry G is spontaneously broken by the F-term superpotential which is holomorphic function of the chiral superfields. As a result, the symmetry G is enhanced to its complexification $G^{\mathbf{C}}$, and the isotropy group H is extended to $\hat{H}$ which contains the complexification $H^{\mathbf{C}}$ of H [5, 3]. BKMU constructed the following three types of Kähler potentials for the target manifold $G^{\mathbf{C}}/\hat{H}$. The A- and C-type Lagrangians contain arbitrary functions of the G-invariant quantities. The B-type Lagrangian is a generalization of the Zumino's Lagrangian [2], and does not have such arbitrariness.

When there is no QNG boson, it has been proved by Itoh, Kugo and Kunitomo (IKK) [4] that the most general Lagrangian is uniquely determined by the B-type

Lagrangian. In this case the target manifold $G^{\mathbf{C}}/\hat{H}$ is isomorphic to G/H, and the unbroken group H must be the centralizer of a torus. This means that the homogeneous manifold G/H should be Kähler. This condition restricts the breaking pattern and is used to classify all the compact homogeneous Kähler G/H Lagrangian [4].

For the low-energy effective Lagrangian of linear field theory origin, there must be at least one QNG boson [5, 7], so that we cannot use the IKK construction. Instead, we must use the A- or C-type Lagrangians of BKMU.[1] As we already remarked, these contain arbitrary functions due to the existence of QNG bosons. The origin of these functions is understood geometrically as follows.

The target manifold $G^{\mathbf{C}}/\hat{H}$ is non-compact and the QNG bosons correspond to the non-compact directions [5, 6, 7]. Since the symmetry of the Lagrangian is still G but not $G^{\mathbf{C}}$, the metric in the QNG directions is not determined by the symmetry, whereas that in the NG directions is determined by the low energy theorems. The arbitrary function thus describes the shape of non-compact directions of the target manifold [6, 7].

The arbitrariness discussed above suggests that the scattering amplitudes of the QNG bosons will not be determined by symmetries and there may not be any low energy theorem. (PCAC relation in supersymmetric theory has been discussed in ref. [8].) In this paper, we investigate the low energy theorems for the scattering amplitudes of NG and QNG bosons, and show that the scattering amplitudes of only the QNG bosons actually coincide with those of NG bosons, in spite of the arbitrariness. The arbitrariness shows up in the scattering amplitudes between NG and QNG bosons. We also confirm that the amplitudes of NG in supersymmetric theories are the same as those in the non-supersymmetric theory.

In sect. 2, we review the CCWZ bosonic theories and the low energy theorems. In sect. 3, we proceed to the supersymmetric theories and derive the low energy theorems. Sect. 4 is devoted to conclusions and discussions.

---

[1] In this paper, we use only the A-type Lagrangian, because little is known about the C-type Lagrangian of BKMU.


$\phi_0^i + \pi^i = \lambda^i(1)$
$\xi^i = \dot{\lambda}^i(0)$
$\lambda^i(t)$
$\phi_0^i = \lambda^i(0)$

Figure 1. Riemann normal coordinate

2 Scattering amplitudes of NG bosons in non-supersymmetric theory

In this section, we start with the bosonic theories. The most general low-energy effective Lagrangian is expressed as

$$\mathcal{L}(\phi) = \frac{1}{2} g_{\alpha\beta}(\phi) \partial_\mu \phi^\alpha \partial^\mu \phi^\beta, \tag{2.1}$$

where we use $\alpha, \beta, \cdots$ for curved indices on the target manifold and μ for spacetime coordinates. This is a nonlinear sigma model on a general target manifold with a metric $g_{\alpha\beta}(\phi)$. The scattering amplitudes calculated from this Lagrangian are invariant under the field redefinitions which may be interpreted as the general coordinate transformations on the target manifold. Using this freedom, we may choose the most convenient Riemann normal coordinate (figure 1). It is the coordinate spanned by the geodesic lines $\lambda^i(t)(0 \leq t \leq 1)$ starting from a point determined by the vacuum expectation value.[2]

To compute scattering amplitudes, we expand the effective Lagrangian around the constant vacuum expectation value ϕ_0 $(\partial_\mu \phi_0 = 0)$ in the Riemann normal coordinate. Since the metric $g_{ij}(\phi_0)$ is diagonal in this coordinate, we denote it as $\frac{1}{2} f_\pi^2 \delta_{ij}$. We find [9]

$$\mathcal{L}(\phi_0 + \pi) = \frac{1}{2} g_{ij}(\phi_0 + \pi) \partial_\mu (\phi_0 + \pi)^i \partial^\mu (\phi_0 + \pi)^j$$

---

[2] We use $i, j, \cdots$ for the flat tangent space indices, which coincide with the indices of broken generators at $\phi = \phi_0$.



Figure 2. Two-body scattering of NG bosons

$$= \frac{1}{2} f_\pi^2 \delta_{ij} \partial_\mu \xi^i \partial^\mu \xi^j + \frac{1}{6} R_{iklj}(\phi_0) \xi^k \xi^l \partial_\mu \xi^i \partial^\mu \xi^j + O(\xi^5), \qquad (2.2)$$

where terms up to the fourth order in ξ^i are kept because they are sufficient in order to compute two-body scattering amplitudes (figure 2). The fluctuating field ξ^i is interpreted as the particle propagating on the vacuum ϕ_0. To calculate many-body scattering amplitudes, we must expand the Lagrangian to higher orders in ξ^i. Using the definition

$$\begin{aligned} &< \xi^k(p_k), \xi^l(p_l) | i\mathcal{L}_{int} | \xi^i(p_i), \xi^j(p_j) > \\ &= i(2\pi)^4 \delta^{(4)}(p_k + p_l - p_i - p_j) \mathcal{M}(\xi^i(p_i), \xi^j(p_j) \to \xi^k(p_k), \xi^l(p_l)), \end{aligned} \qquad (2.3)$$

we find that the two-body scattering amplitudes are in general expressed in terms of the curvature tensor as [3]

$$\begin{aligned} &\mathcal{M}(\xi^i(p_i), \xi^j(p_j) \to \xi^k(p_k), \xi^l(p_l)) \\ &= -\frac{1}{3f_\pi^4}[(s-u)R_{kijl} + (u-t)R_{kjil} + (t-s)R_{kjli}], \end{aligned} \qquad (2.4)$$

where the Madelstam variables s, t and u are defined by

$$\begin{aligned} s &\overset{\text{def}}{=} (p_i + p_j)^2 = +2p_i \cdot p_j = +2p_l \cdot p_k, \\ t &\overset{\text{def}}{=} (p_i - p_k)^2 = -2p_i \cdot p_k = -2p_l \cdot p_j, \\ u &\overset{\text{def}}{=} (p_i - p_l)^2 = -2p_i \cdot p_l = -2p_j \cdot p_l. \end{aligned} \qquad (2.5)$$

---

[3] Since two point function of ξ is proportional to f_π^{-2}, we have to supply f_π^{-4} to obtain the scattering amplitude properly normalized.

Our task is now to determine the metric for the target manifold of the effective Lagrangian of NG bosons by the symmetries G and H. The coset manifold G/H is parametrized by the NG bosons ϕ^α. We denote the broken and unbroken Hermitian generators by[4]

$$X_i \in \mathcal{G} - \mathcal{H}\ ,\ \ H_a \in \mathcal{H}. \tag{2.6}$$

The fundamental variable in the construction of the metric is the representative of the coset G/H

$$U(\phi) = e^{i\phi\cdot X} \in G/H\ ,\ \ \phi\cdot X = \phi^\alpha X_i \delta^i_\alpha, \tag{2.7}$$

which is unitary.

In order to express the metric in terms of the fundamental variable, it is useful to introduce the Maurer-Cartan 1-form

$$\frac{1}{i}U^{-1}dU = (e^i_\alpha(\phi)X_i + \omega^a_\alpha(\phi)H_a)d\phi^\alpha, \tag{2.8}$$

which is a Lie algebra valued 1-form. The broken generator element of the Maurer-Cartan 1-form, $e^i_\alpha(\phi)$ is called the vielbein, which connects the metric of target manifold and the tangent space flat metric. The unbroken element $\omega^a_\alpha(\phi)$ is called the canonical H connection. Using the definition given in (2.8), we can explicitly write the metric of the target manifold

$$g_{\alpha\beta}(\phi) = f_\pi^2 e^i_\alpha(\phi)e^j_\beta(\phi)\delta_{ij}, \tag{2.9}$$

by the decay constant of the pion (NG boson) f_π, which has unit mass dimension and represents the size of the coset manifold G/H. If G/H is reducible, there are as many decay constants as the number of the H irreducible factors; if it is irreducible, there exists only one decay constant.

The Maurer-Cartan's structure equation

$$d(U^{-1}dU) + (U^{-1}dU)\wedge(U^{-1}dU) = 0, \tag{2.10}$$

---

[4] The Lie algebras of the groups G and H are denoted by $\mathcal{G}$ and $\mathcal{H}$, respectively. We use the indices $i, j, \cdots$ for the broken generators and $a, b, \cdots$ for the unbroken generators.

enables us to calculate the curvature tensor of the symmetric irreducible coset manifold G/H as [7]

$$R_{ijkl} = f_\pi^2 \, f_{ij}{}^a f_{akl}. \tag{2.11}$$

The low energy theorems for the two-body scattering amplitudes of NG bosons then take the form

$$\begin{aligned} &\mathcal{M}(\xi^i(p_i), \xi^j(p_j) \to \xi^k(p_k), \xi^l(p_l)) \\ &= -\frac{1}{3f_\pi^2}[(s-u)f_{ki}{}^a f_{ajl} + (u-t)f_{kl}{}^a f_{aij} + (t-s)f_{kj}{}^a f_{ali}]. \end{aligned} \tag{2.12}$$

This is a well-known result in the current algebra [10]. On the other hand, eq. (2.4) is valid in more general cases. If the target manifold G/H is non-symmetric or reducible, the curvature tensor is more complicated [7] and so are the low energy theorems.

In conclusion, the scattering amplitudes are written by the geometrical quantities (for example, the curvature tensor in the case of the two-body scattering amplitudes). The symmetries restrict the form of the metric tensor of the target manifold of the sigma model and are strong enough to determine the geometric objects (curvature tensor) in terms of group structure constants up to an overall factor.

In the next section, we introduce supersymetry in addition to the global symmetries G and H. At first glance, the restriction seems to be stronger and the low energy theorems more restrictive. We will see that this is not true because supersymmetry requires extra massless bosons and the target manifolds become more complicated.

3 Scattering amplitudes of NG and QNG bosons in supersymmetric theory

In $N = 1$ supersymmetric theories, the general Lagrangian for chiral fileds is

$$\mathcal{L} \;=\; \int d^2\theta d^2\bar{\theta} K(\Phi, \Phi^\dagger)$$

$$
\begin{aligned}
&= g_{ij^*}(\varphi,\varphi^*)\partial_\mu\varphi^i\partial^\mu\varphi^{*j} + ig_{ij^*}\bar{\psi}^j\bar{\sigma}^\mu(\partial_\mu\psi^i + \Gamma^i{}_{lk}\partial_\mu\varphi^l\psi^k) \\
&\quad + \frac{1}{4}R_{ij^*kl^*}\psi^i\psi^k\bar{\psi}^j\bar{\psi}^l,
\end{aligned} \tag{3.1}
$$

where the metric tensor of the target Kähler manifold [11] is given by the Kähler potential K as[5]

$$
g_{ij^*}(\varphi,\varphi^*) = \frac{\partial^2 K(\varphi,\varphi^*)}{\partial\varphi^i\partial\varphi^{*j}}. \tag{3.2}
$$

The chiral superfield has the component fields

$$
\Phi = \varphi + \sqrt{2}\theta\psi + \theta\theta F. \tag{3.3}
$$

The bosonic part is a complex scalar field

$$
\varphi = A + iB, \tag{3.4}
$$

which is the coordinate of the target manifold. Here A bosons always represent the ordinary NG bosons, but B bosons may be NG bosons or QNG bosons. In the former case the chiral superfields are called the pure type or the non-doubled type; in the latter case they are called the mixed type or the doubled type. The maximal realization is the nonlinear realization in which A bosons are NG bosons but all B bosons are QNG bosons. In this paper we focus on this maximal realization because it is the most natural one in Lagrangian field theories. Although we do not discuss the fermions in this paper, fermions in the chiral superfields are sometimes called the quasi Nambu-Goldstone fermions.

For the low-energy effective Lagrangian with the symmetry G broken down to H, the Kähler potential must have G-isometry invariance (up to the Kähler transformation). BKMU constructed G-invariant Kähler potential by generalizing the CCWZ method. The broken and unbroken generators of the complexified group are written as

$$
Z_R \in \mathcal{G}^{\mathbf{C}} - \hat{\mathcal{H}}\ ,\ K_M \in \hat{\mathcal{H}}. \tag{3.5}
$$

---

[5] Here we use the indices $i, j, \cdots$ for general complex coordinates of the Kähler target manifold. They should not be confused with the indices of broken generators in non-supersymmetric case.

The commutators of these generators are given by

$$[K_M, K_N] = i f_{MN}{}^L K_L \ , \ [Z_R, K_M] = i f_{RM}{}^N K_N \ , \ [Z_R, Z_S] = i f_{RS}{}^T Z_T, \tag{3.6}$$

where in the last equation, we have assumed that the complex coset $G^{\mathbf{C}}/\hat{H}$ is a symmetric space with an involutive automorphism

$$Z_R \to -Z_R \ , \ K_M \to K_M. \tag{3.7}$$

In general, these generators are written by complex linear combinations of the Hermitian generators of G. The Hermitian generators in $\mathcal{G}^{\mathbf{C}} - \hat{\mathcal{H}}$ correspond to the mixed type chiral superfileds, while the sets of non-Hermitian step generators in $\mathcal{G}^{\mathbf{C}} - \hat{\mathcal{H}}$ and $\hat{\mathcal{H}}$ correspond to the pure type ones.

A novel feature in the supersymmetric theories is that the BKMU fundamental variable

$$\xi(\Phi) = \mathrm{e}^{i\Phi \cdot Z} \in G^{\mathbf{C}}/\hat{H} \ , \ \Phi \cdot Z = \sum_{i=1}^{N_\Phi} \Phi^i Z_R \delta_i^R, \tag{3.8}$$

is not unitary. This is due to the complex coordinates (3.4). This variable is the representative of the complex coset $G^{\mathbf{C}}/\hat{H}$. The transformation law of ξ is

$$\xi \xrightarrow{g} \xi' = g\xi\hat{h}^{-1}(g,\xi). \tag{3.9}$$

In order to write down Kähler potential, let us now prepare some representation (ρ, V) of G. In the case of low-energy effective Lagrangian of linear model origin, there are some $\hat{H}$-invariant vectors $v_a, v_b, \cdots \in V$ in the representation space:

$$\rho(\hat{H})v_a = v_a. \tag{3.10}$$

The G-invariant Kähler potential (BKMU A-type Lagrangian) is then an arbitrary function of G-invariant quantities [5, 3]

$$K(\Phi, \Phi^\dagger) = f(v_a^\dagger \rho(\xi^\dagger(\Phi^\dagger)\xi(\Phi))v_b). \tag{3.11}$$

For simplicity, we assume that there is only one $\hat{H}$ invariant vector v in the representation space. In this case the coset space G/H is a symmetric one [5, 7]. This assumption is not so restrictive: For example, this admits the case of the chiral symmetry breaking, where $G = G_L \times G_R$ breaks down to $H = G_V$ and the full target manifold is $G_L^{\mathbf{C}} \times G_R^{\mathbf{C}}/G_V^{\mathbf{C}}$ [3].

In supersymmetric theories, it is useful to use the complex coset $G^{\mathbf{C}}/\hat{H}$ version of the Maurer-Cartan 1-form (2.8):

$$\frac{1}{i}\xi(\varphi)^{-1}d\xi(\varphi) = (E_i^R(\varphi)Z_R + W_i^M(\varphi)K_M)d\varphi^i. \tag{3.12}$$

The broken generator element of the Maurer-Cartan 1-form $E_i^R(\varphi)$ is called the holomorphic vielbein [7]. This is different from the ordinary non-holomorphic vielbein connecting the Kähler and tangent-space flat metrics. The unbroken generator element $W_i^M(\varphi)$ is called the canonical $\hat{H}$ connection, which, together with a Kähler potential, fixes the spin connection in the holomorphic formalism [6, 7]. These can be written in the coordinate φ explicitly as

$$\begin{aligned} E_i^R(\varphi) &= i\delta_i^S\left(\frac{1-e^{i\varphi\cdot T}}{\varphi\cdot T}\right)_S^{\ \ R} \\ &= \delta_i^S\left(1+\frac{1}{2!}i\varphi\cdot T+\frac{1}{3!}(i\varphi\cdot T)^2+\cdots\right)_S^{\ \ R}, \qquad (3.13) \\ W_i^M(\varphi) &= i\delta_i^S\left(\frac{1-e^{i\varphi\cdot T}}{\varphi\cdot T}\right)_S^{\ \ M} \\ &= \delta_i^S\left(1+\frac{1}{2!}i\varphi\cdot T+\frac{1}{3!}(i\varphi\cdot T)^2+\cdots\right)_S^{\ \ M}. \qquad (3.14) \end{aligned}$$

Here T is the generator of the adjoint representation of $\mathcal{G}$; $(T_C)_A{}^B = if_{AC}{}^B$. We can always choose the point $\varphi = 0$ to be a symmetric point, where all QNG bosons are set to zero, since any symmetric points are equivalent under the G isometry. At this point $\varphi = 0$, these reduce to

$$E_i^R|_{\varphi=0} = \delta_i^R, \tag{3.15}$$

$$W_i^M|_{\varphi=0} = 0. \tag{3.16}$$

For later convenience, let us calculate the derivatives of these at the point $\varphi = 0$:

$$\partial_j E_i^R|_{\varphi=0} = -\frac{1}{2}\delta_i^S\delta_j^T f_{ST}{}^R = 0, \tag{3.17}$$

$$\partial_j W_i^M|_{\varphi=0} = -\frac{1}{2}\delta_i^S\delta_j^T f_{ST}{}^M. \tag{3.18}$$

In the first equation, we have assumed that $G^{\mathbf{C}}/\hat{H}$ is symmetric.

To calculate geometrical quantities in the holomorphic formalism, the following formulas resulting from (3.12) are useful:

$$\begin{aligned}
\frac{\partial \xi}{\partial \varphi^i} v &= i\xi E_i^R Z_R v, \\
\frac{\partial \xi}{\partial \varphi^i} Z_S v &= i\xi (E_i^R Z_R Z_S + W_i^M [K_M, Z_S]) v \\
&= i\xi (E_i^R Z_R Z_S + i W_i^M f_{MS}{}^T Z_T) v. \qquad (3.19)
\end{aligned}$$

Using these formulas, we calculate the metric tensor from (3.11)

$$g_{ij^*}(\varphi, \varphi^*) = G_{RS^*}(\varphi, \varphi^*) E_i^R(\varphi) (E_j^S(\varphi))^*, \qquad (3.20)$$

where

$$G_{RS^*} = f'(v^\dagger \xi^\dagger \xi v)(v^\dagger Z_S^\dagger \xi^\dagger \xi Z_R v) + f''(v^\dagger \xi^\dagger \xi v)(v^\dagger Z_S^\dagger \xi^\dagger \xi v)(v^\dagger \xi^\dagger \xi Z_R v), \qquad (3.21)$$

is an auxiliary metric [7]. Note that this is not equal to the Hermitian metric of the (flat) tangent space.

The conventional G/H sigma model is embedded in the full manifold $G^{\mathbf{C}}/\hat{H}$ [7], and the coordinates are spanned by QNG as well as NG bosons in the form of (3.4). If there are some pure type multiplets, the corresponding B fields are NG bosons and the BKMU (3.8) and CCWZ (2.7) variables do not coincide even if we put all QNG bosons to zero; we need a local $\hat{H}$ transformation from the right to compensate the difference [4, 6, 7]. In the maximal realization we are considering, however, the imaginary directions are all given by QNG bosons and so the CCWZ (2.7) and BKMU (3.8) variables are simply connected by

$$\xi(\varphi)|_{\mathrm{QNG}=0} = U(\phi). \qquad (3.22)$$

This means that at the symmetric point, $\xi^\dagger \xi = 1$ and (3.21) reduces to

$$G^0_{RS^*} = f'(v^\dagger v)(v^\dagger Z_S^\dagger Z_R v) + f''(v^\dagger v)(v^\dagger Z_S^\dagger v)(v^\dagger Z_R v), \qquad (3.23)$$

where the superscript 0 stands for quantities at the symmetric point (QNG=0).

The symmetric space $G^{\mathbf{C}}/\hat{H}$ is characterized by the involutive automorphism (3.7). Since $\hat{H}$ invariant vector v can be chosen as an eigenvector of the involution with eigenvalues ±1, $v^\dagger Z_R v$ changes sign under the involution and hence vanishes. By a suitable choice of broken generators, $Z_S v$ becomes orthonormal basis of the representation space and we have $v^\dagger Z_R^\dagger Z_S v = \delta_{R^* S}$. The auxiliary metric at the symmetric point then becomes

$$\begin{aligned} G^0_{RS^*} &= f'(v^\dagger v) v^\dagger v \ \delta_{RS^*} \\ &= \frac{1}{2} f_\pi^2 \delta_{RS^*}, \end{aligned} \tag{3.24}$$

where we have defined the decay constant of the pion (NG boson) by

$$f_\pi^2 \overset{\text{def}}{=} 2 f'(v^\dagger v) v^\dagger v, \tag{3.25}$$

which characterizes the size of the compact submanifold G/H. The normalization in (3.25) is fixed so as to give canonically normalized kinetic terms for NG bosons.

For the maximal realization of our interest, the complex version of the Maurer-Cartan form (3.12) becomes the conventional one (2.8) without any local $\hat{H}$ transformation because of the relation (3.22). So the holomorphic vielbein becomes the ordinary vielbein on G/H, and the effective Lagrangian for A bosons takes the same form as the bosonic case:

$$\mathcal{L}_{\text{boson}}|_{\text{QNG}=0} = \frac{1}{2} f_\pi^2 e^i_\alpha e^j_\beta \delta_{ij} \partial_\mu A^\alpha \partial^\mu A^\beta = \frac{1}{2} g_{\alpha\beta} \partial_\mu A^\alpha \partial^\mu A^\beta. \tag{3.26}$$

The fact that we recover the conventional G/H model by putting QNG$= 0$ in the effective Lagrangian can be understood naturally as follows: If supersymmetry is broken, the QNG bosons get the mass of the supersymmetry breaking scale. We can then integrate out the massive QNG bosons, and obtain the conventional G/H model, making the curvature tensor of the obtained manifold identical to eq. (2.11). The low energy theorems for the scattering amplitudes of NG bosons are the same as eq. (2.12).

From now on we concentrate on the geometric quantities at the symmetric point. We apply these to the low-energy scattering amplitudes of NG and QNG bosons. Let us first show that the coset coordinate φ is a special one called the adapted coordinate. According to eqs. (3.15) and (3.24), the ordinary metric at the point $\varphi = 0$ becomes

$$g_{ij^*}|_{\varphi=0} = G^0_{RS^*} E^R_i|_{\varphi=0} (E^S_j)^*|_{\varphi^*=0} = \frac{1}{2} f_\pi^2 \delta_{ij^*}. \tag{3.27}$$

The derivative of the auxiliary metric is

$$\begin{aligned}
\partial_j G_{RS^*} = {} & i \Big\{ f'(z)(v^\dagger Z_S^\dagger \xi^\dagger \xi Z_T Z_R v) + f''(z)(v^\dagger \xi^\dagger \xi Z_T v)(v^\dagger Z_S^\dagger \xi^\dagger \xi Z_R v) \\
& + f''(z)(v^\dagger Z_S^\dagger \xi^\dagger \xi Z_T v)(v^\dagger \xi^\dagger \xi Z_R v) + f''(z)(v^\dagger Z_S^\dagger \xi^\dagger \xi v)(v^\dagger \xi^\dagger \xi Z_T Z_R v) \\
& + f'''(z)(v^\dagger \xi^\dagger \xi Z_T v)(v^\dagger Z_S^\dagger \xi^\dagger \xi v)(v^\dagger \xi^\dagger \xi Z_R v) \Big\} E^T_j \\
& + i \Big\{ f'(z)(v^\dagger Z_S^\dagger \xi^\dagger \xi Z_T v) + f''(z)(v^\dagger Z_S^\dagger \xi^\dagger \xi v)(v^\dagger \xi^\dagger \xi Z_T v) \Big\} i f_{MR}{}^T W^M_j,
\end{aligned} \tag{3.28}$$

where $z = v^\dagger \xi^\dagger \xi v$. The derivative of the metric (3.20) is

$$\partial_j g_{kl^*} = (\partial_j G_{RS^*}) E^R_k (E^S_l)^* + G_{RS^*} (\partial_j E^R_k)(E^S_l)^*. \tag{3.29}$$

At the symmetric point, this reduces to

$$\partial_j g_{kl^*}{}^0 = \frac{1}{2} f_\pi^2 (-f_{MRS^*} W^M_j E^R_k (E^S_l)^* + \delta_{RS^*} (\partial_j E^R_k)(E^S_l)^*)^0, \tag{3.30}$$

and especially at the point $\varphi = 0$ to

$$\partial_j g_{kl^*}|_{\varphi=0} = 0, \tag{3.31}$$

as a result of eqs. (3.16) and (3.17) when $G^{\mathbf{C}}/\hat{H}$ is symmetric. Since the inverse metric at the symmetric point is

$$g^{ij^*0} = 2 f_\pi^{-2} \delta^{RS^*} (E^i_R (E^j_S)^*)^0, \tag{3.32}$$

the (coordinate) connection at the symmetric point is given by

$$\begin{aligned}
\Gamma^i{}_{jk}{}^0 &= (g^{il^*} \partial_j g_{kl^*})^0 \\
&= (-f_{MR}{}^S W^M_j E^R_k E^i_S + (\partial_j E^S_k) E^i_S)^0,
\end{aligned} \tag{3.33}$$

and especially on the point $\varphi = 0$ by

$$\Gamma^i{}_{jk}|_{\varphi=0} = 0 \tag{3.34}$$

due to eqs. (3.16) and (3.17) when $G^{\mathbf{C}}/\hat{H}$ is symmetric. The coordinate with the properties (3.27) and (3.34) is called the adapted coordinate on the point $\varphi = 0$.[6]

Because of eqs. (3.31), (3.17) and (3.15), the curvature tensor in the adapted coordinate becomes

$$\begin{aligned} R_{ij^*kl^*}|_{\varphi=0} &= g_{ij^*,kl^*}|_{\varphi=0} = \frac{\partial^4 K}{\partial\varphi^i\partial\varphi^{*j}\partial\varphi^k\partial\varphi^{*l}}|_{\varphi=0} \\ &= (\partial_k\partial_{l^*}G_{RS^*})|_{\varphi=0}\delta_i^R(\delta_j^S)^*. \end{aligned} \tag{3.35}$$

According to eq. (3.28), the second derivative of the auxiliary metric is

$$\begin{aligned} (\partial_k\partial_{l^*}G_{RS^*})|_{\varphi=0} &= [f'(v^\dagger v)(v^\dagger Z_S^\dagger Z_V^\dagger Z_U Z_R v) \\ &\quad + g^2(\delta_{RU}\delta_{S^*V^*} + \delta_{RV^*}\delta_{US^*} + \delta_{RS^*}\delta_{UV^*})]\delta_k^U(\delta_l^V)^*, \end{aligned} \tag{3.36}$$

and the curvature tensor at the point $\varphi = 0$ in the adapted coordinate is

$$\begin{aligned} R_{ij^*kl^*}|_{\varphi=0} &= [f'(v^\dagger v)(v^\dagger Z_S^\dagger Z_V^\dagger Z_U Z_R v) + g^2(\delta_{RU}\delta_{S^*V^*} + \delta_{RV^*}\delta_{US^*} + \delta_{RS^*}\delta_{UV^*})] \\ &\quad \times\delta_i^R(\delta_j^S)^*\delta_k^U(\delta_l^V)^*, \end{aligned} \tag{3.37}$$

where we have defined the constant g by

$$g^2 \overset{\text{def}}{=} f''(v^\dagger v)(v^\dagger v)^2. \tag{3.38}$$

The constant f_π determines the total size of the compact submanifold G/H. On the other hand, the constant g and higher derivatives of the arbitrary function f control the shape of the non-compact directions of the target manifold. Note that the curvature (3.37) has the symmetry of Kähler manifold $R_{ij^*kl^*} = R_{kj^*il^*}$, since $R_{ij^*kl^*} - R_{kj^*il^*}$ contains the commutators of broken generators, which produce unbroken generators when $G^{\mathbf{C}}/\hat{H}$ is symmetric, and vanishes upon multiplying the $\hat{H}$ invariant vector v.

In sect. 2, we have found that the scattering amplitudes are written in terms of the curvature tensors of the target manifold. Let us repeat the same analysis here with the above formalism. Supersymmetry requires that the target manifold should be Kähler. This requirement restricts the form of the curvature tensor of the target manifold and also the low energy theorems. The low-energy effective Lagrangian is expanded by the fields φ and ψ around the point $\varphi = 0$ up to fourth terms as

---

[6] That we can choose an adapted coordinate on any point is a necessary and sufficient condition that a complex manifold should be a Kähler manifold.

$$\begin{aligned} \mathcal{L} &= g_{ij^*}|_{\varphi=0}\, \partial_\mu \varphi^i \partial^\mu \varphi^{*j} + i g_{ij^*}|_{\varphi=0}\, \bar{\psi}^j \bar{\sigma}^\mu \partial_\mu \psi^i \\ &+ R_{ij^*kl^*}|_{\varphi=0}\, \varphi^k \varphi^{*l} \partial_\mu \varphi^i \partial^\mu \varphi^{*j} + \frac{1}{4} R_{ij^*kl^*}|_{\varphi=0}\, \psi^i \psi^k \bar{\psi}^j \bar{\psi}^l \\ &+ i R_{ij^*kl^*}|_{\varphi=0}\, \varphi^{*j} \partial_\mu \varphi^i (\bar{\psi}^l \bar{\sigma}^\mu \psi^k). \end{aligned} \tag{3.39}$$

In deriving this result, we have made a holomorphic coordinate transformation to eliminate non-covariant terms. The last three terms are the interaction Lagrangians, which are described by the curvature tensor of the Kähler manifold. They give the two-body scattering amplitudes by the curvature.

The low-energy scattering amplitudes are directly connected with the curvature tensor in the real basis. In such basis, the components of the curvature tensor in the Kähler manifold are divided into four types

$$\begin{cases} R_{A^iA^jA^kA^l} = R_{B^iB^jB^kB^l} = R_{A^iA^jB^kB^l} = R_{B^iB^jA^kA^l} \\ R_{B^iA^jB^kA^l} = R_{A^iB^jA^kB^l} = -R_{B^iA^jA^kB^l} = -R_{A^iB^jB^kA^l} \\ R_{B^iA^jA^kA^l} = -R_{A^iB^jA^kA^l} = -R_{A^iB^jB^kB^l} = R_{B^iA^jB^kB^l} \\ R_{A^iA^jB^kA^l} = -R_{A^iA^jA^kB^l} = -R_{B^iB^jA^kB^l} = R_{B^iB^jB^kA^l} \end{cases}, \tag{3.40}$$

as a result of the Kähler conditions.[7] These curvatures have the symmetry under

$$\begin{cases} A^i \to +B^i \\ B^i \to -A^i \end{cases}, \tag{3.41}$$

and so do the scattering amplitudes.

The first equation in (3.40) implies that the two-body amplitudes of the QNG bosons coincide with those of the corresponding NG bosons

$$\mathcal{M}(\mathrm{NG}, \mathrm{NG} \to \mathrm{NG}, \mathrm{NG}) = \mathcal{M}(\mathrm{QNG}, \mathrm{QNG} \to \mathrm{QNG}, \mathrm{QNG}), \tag{3.42}$$

owing to eq. (2.4) (see figure 3). Note that this is a result following solely from the Kähler conditions and is valid for general Kähler sigma models.

---

[7] These are equivalent to the vanishing curvature conditions $R_{ijkl} = R_{i^*jkl} = R_{ij^*kl} = R_{ijk^*l} = R_{ijkl^*} = R_{i^*j^*kl} = 0$ and their Hermitian conjugates.



Figure 3. Scattering amplitudes of QNG coincide with those of NG

Let us now evaluate the actual values of the curvature tensors in the real basis. By using (3.37) and

$$\partial_{A^i} = \partial_i + \partial_i^* \ , \ \partial_{B^i} = i(\partial_i - \partial_i^*), \tag{3.43}$$

we derive[8]

$$\begin{aligned} R_{A^iA^jA^kA^l}|_{\varphi=0} &= R_{ij^*kl^*} + R_{i^*jkl^*} + R_{ij^*k^*l} + R_{i^*jk^*l} \\ &= R_{ij^*kl^*} - R_{ji^*kl^*} - R_{ij^*lk^*} + R_{ji^*lk^*} \\ &= f'(v^\dagger v) v^\dagger (Z_S[Z_V, Z_U]Z_R - Z_R[Z_V, Z_U]Z_S) v \delta_i^R \delta_j^S \delta_k^U \delta_l^V \\ &= f_\pi^2 f_{RS}{}^M f_{MUV} \delta_i^R \delta_j^S \delta_k^U \delta_l^V , \end{aligned} \tag{3.44}$$

where we have also used eq. (3.25). This depends only on the first derivative of the arbitrary function and the higher derivatives all cancel out. It is thus completely independent of the shape of the non-compact directions. This result demonstrates that the curvature tensors and hence the low energy theorems for the NG bosons coincide with those in non-supersymmetric theories.

Similarly, we can obtain the components of the curvature tensor involving both the NG and QNG bosons. One finds

$$\begin{aligned} R_{B^iA^jB^kA^l}|_{\varphi=0} &= i^2 (R_{ij^*kl^*} - R_{i^*jkl^*} - R_{ij^*k^*l} + R_{i^*jk^*l}) \\ &= -f'(v^\dagger v) v^\dagger (Z_S\{Z_V, Z_U\}Z_R + Z_R\{Z_V, Z_U\}Z_S) v \delta_i^R \delta_j^S \delta_k^U \delta_l^V \\ &\quad -4g^2(\delta_{ik}\delta_{jl} + \delta_{il}\delta_{kj} + \delta_{ij}\delta_{kl}). \end{aligned} \tag{3.45}$$

---

[8] For notational simplicity, we do not write $\varphi = 0$ on the right hand side.

These components include higher derivatives of the arbitrary function, which controls the non-compact QNG directions, showing their dependence on the shape of non-compact directions and the details of the underlying fundamental theory. The low-energy scattering amplitudes for two NG bosons and two QNG bosons are calculated by use of eqs. (3.45), (3.40) and (2.4). They also depend on the higher derivatives of the arbitrary function.

We also find from (3.40)[9]

$$\begin{aligned} R_{B^iA^jA^kA^l}|_{\varphi=0} &= i(R_{ij^*kl^*} - R_{i^*jkl^*} + R_{ij^*k^*l} - R_{i^*jk^*l}) \\ &= if'(v^\dagger v)v^\dagger(Z_S[Z_V, Z_U]Z_R + Z_R[Z_V, Z_U]Z_S)v\delta_i^R\delta_j^S\delta_k^U\delta_l^V \\ &= 0, \end{aligned} \tag{3.46}$$

$$\begin{aligned} R_{A^iA^jB^kA^l}|_{\varphi=0} &= i(R_{ij^*kl^*} + R_{i^*jkl^*} - R_{ij^*k^*l} - R_{i^*jk^*l}) \\ &= if'(v^\dagger v)v^\dagger(Z_V[Z_S, Z_R]Z_U + Z_U[Z_S, Z_R]Z_V)\delta_i^R\delta_j^S\delta_k^U\delta_l^V \\ &= 0. \end{aligned} \tag{3.47}$$

From these two equations and eq. (3.40), we find that all components of the curvature tensor which include odd numbers of NG bosons and QNG bosons vanish. Eq. (2.4) then tells us that the low-energy scattering amplitudes with odd numbers of NG bosons and QNG bosons all vanish.

4 Conclusions and Discussions

In supersymmetric theories whose global symmetry G is broken down to its subgroup H, there appear the QNG bosons besides the ordinary NG bosons. The low-energy effective Lagrangian can be described by a nonlinear sigma model whose target manifold is a Kähler coset $G^{\mathbf{C}}/\hat{H}$ with only the compact symmetry G. The manifold is non-compact and its non-compact directions correspond to the QNG bosons. Since the global symmetry does not restrict the shape of the non-compact directions, the Kähler potential of the effective Lagrangian includes an arbitrary function.

---

[9] Actually these two quantities in (3.46) and (3.47) are not independent since $R_{B^iA^jA^kA^l} = R_{A^kA^lB^iA^j}$.

Nevertheless we have shown that the low-energy scattering amplitudes which has only QNG bosons are equal to those of the corresponding NG bosons. The arbitrariness comes into the low-energy scattering amplitudes which has two NG bosons and two QNG bosons. The low-energy scattering amplitudes which have the odd number of NG bosons and QNG bosons all vanish for the case of symmetric point.

In this article, we have investigated only the two-body scattering amplitudes. However, generalization to many-body scattering amplitudes is straightforward. If we expand the Lagrangian in the Riemann normal coordinate to higher orders in ξ, the coefficients are covariant geometrical objects such as covariant derivatives of curvature tensor. We can obtain (exact) low-energy scattering amplitudes by using the effective Lagrangian. They will have the symmetry NG $\rightarrow$ QNG, QNG $\rightarrow -$NG similar to the two-body amplitudes, and we can obtain the low energy theorems of NG and QNG bosons.

Another interesting extension of our work is to examine how our results are generalized at non-symmetric point. We expect that such a generalization brings new features into the low energy theorems.

Acknowledgement

One of the authors (M. Nitta) is very grateful to S. Iso and N. Ishibashi for useful discussions.

References

[1] S. Coleman, J. Wess and B. Zumino, Phys. Rev. **177** (1969) 2239;
C. G. Callan, S. Coleman, J. Wess and B. Zumino, Phys. Rev. **177** (1969) 2247.

[2] B. Zumino, Phys. Lett. **87B** (1979) 203.

[3] M. Bando, T. Kuramoto, T. Maskawa and S. Uehara, Phys. Lett. **138B** (1984) 94; Prog. Theor. Phys. **72** (1984) 313, 1207.

[4] K. Itoh, T. Kugo and H. Kunitomo, Nucl. Phys. **B263** (1986) 295; Prog. Theor. Phys. **75** (1986) 386.

[5] W. Lerche, Nucl. Phys. **B238** (1984) 582.

[6] W. Buchmüller and W. Lerche, Ann. Phys. **B175** (1987) 159.

[7] G .M . Shore, Nucl. Phys. **B320** (1989) 202; Nucl. Phys. **B334** (1990) 172;
A. C. Kotcheff and G. M. Shore, Int. Jour. Mod. Phys. **A4** (1989) 4391; Nucl. Phys. **B333** (1990) 701.

[8] W. Lerche, Nucl. Phys. **B246** (1984) 475.

[9] L. Alvarez-Gaumé, D. Z. Freedman and S. Mukhi, Ann. Phys. **134** (1981) 85.

[10] M. Bando, T. Kugo and K. Yamawaki, Phys. Rep. **164** (1988) 217.

[11] J. Wess and J. Bagger, *Supersymmetry and Supergravity*, Princeton Univ. Press, Princeton(1992).